

# Occurrence of viruses in Calla and Peruvian lily in Tuscan nurseries and evidence of new viral records in Italy

A. Luvisi <sup>1,2(\*)</sup>, D. Rizzo <sup>3</sup>, L. Stefani <sup>3</sup>, A. Panattoni <sup>2</sup>, A. Materazzi <sup>2</sup>

<sup>1</sup> Dipartimento di Scienze e Tecnologie Biologiche ed Ambientali, Centro Ecotekne Università del Salento, Via Provinciale Lecce Monteroni, 73100 Lecce, Italy.

<sup>2</sup> Dipartimento di Scienze Agrarie, Alimentari e Agro-ambientali, Università di Pisa, Via del Borghetto, 80, 56124 Pisa, Italy.

<sup>3</sup> Servizio Fitosanitario Regionale, Regione Toscana, Via Ciliegiole, 99, 51100 Pistoia, Italy.

**Key words:** *Alstroemeria*, RT-PCR, *Zantedeschia*.

**Abstract:** In order to evaluate the health status of Calla and Peruvian lily in Tuscan nurseries, 18 viruses belonging to six families and one unassigned virus were assayed. Tests were carried out on 90 *Zantedeschia aethiopica* plants and 48 *Alstroemeria* spp. plants collected from 12 Tuscan nurseries in two years, via RT-PCR tests. *Z. aethiopica* was mainly affected by viruses belonging to the *Potyviridae* family, with the main infection caused by Dasheen mosaic virus (DsMV) and *Zantedeschia* mild mosaic virus (ZaMMV). Even if *Alstroemeria* spp. plants were affected by *Potyviridae* family viruses too, higher infection rates were recorded for *Betaflexiviridae*, where Lily symptomless virus infected more than half of plants. This is the first known report of Lily mottle virus (LMoV) in *Alstroemeria* spp. and *Z. aethiopica* or ZaMMV in *Alstroemeria* spp. in Italy.

## 1. Introduction

Calla and Peruvian lily are commonly cultivated in Tuscany (Italy) and they represent one of the main cut flower productions. Calla lily [*Zantedeschia aethiopica* (L.) Spreng] is known to be susceptible to at least 13 virus species, mainly belonging to *Potyviridae*, *Bunyaviridae*, and *Tombusviridae* (Huang *et al.*, 2007). Peruvian lily (*Alstroemeria* spp.) has become one of the most popular cut flowers worldwide. It has been reported as the natural host of various plant viruses, including members of *Potyviridae*, *Betaflexiviridae* or *Bunyaviridae* (Park *et al.*, 2010).

In Tuscany, virus surveys were carried out on various woody plants such as grapevine (Rizzo *et al.*, 2012; 2015 a) but to our knowledge no reports are available for Calla and Peruvian lily in Italy. In order to evaluate the health status of these plants in Tuscan nurseries, various viruses were assayed, belonging to the following families: *Betaflexiviridae* [*Alstroemeria* carla virus (AICV), Lily symptomless

virus (LSV)], *Bromoviridae* [Cucumber mosaic virus (CMV)], *Bunyaviridae* [Impatiens necrotic spot virus (INSV), Iris yellow spot virus (IYSV), Tomato spotted wilt virus (TSWV)], *Comoviridae* [Arabis mosaic virus (ArMV), Broad bean wilt virus 1 (BBWV-1), Broad bean wilt virus 2 (BBWV-2)], *Potyviridae* [*Alstroemeria* mosaic virus (AIMV), Bean yellow mosaic virus (BYMV), Dasheen mosaic virus (DsMV), Konjac mosaic virus (KoMV), Lily mottle virus (LMoV), Turnip mosaic virus (TuMV), *Zantedeschia* mosaic virus (ZaMV), *Zantedeschia* mild mosaic virus (ZaMMV)], *Tombusviridae* [Carnation mottle virus (CarnMV)] and unassigned [Tobacco rattle virus (TRV)]. An additional aim of this survey was to evidence new viral records in Italy for these widespread cultivations, due to the intense international exchanges that characterize the commercialization of lily.

## 2. Materials and Methods

Tests were carried out on 90 *Z. aethiopica* plants and 48 *Alstroemeria* spp. plants collected from 12 Tuscan nurseries in two years. Plants showed symp-

(\*) Corresponding author: andrea.luvisi@unisalento.it  
Received for publication 24 November 2015  
Accepted for publication 22 February 2016

toms such as foliar chlorosis, yellow spot and stripes. Total RNA was extracted from foliar tissue (2 g) using RNeasy Plant Mini Kit (Qiagen, Netherlands) protocol, modified according to MacKenzie *et al.* (1997). Tissues (2 g) were ground using a Tissue lyser (Qiagen) adding 5 ml of grinding buffer (4.0 M guanidine isothiocyanate, 0.2 M sodium acetate pH 5.0, 25 mM EDTA, 2.5% PVP-40 and 2.0% sodium bisulfate) just before use. The homogenate (1 ml) was transferred to a 1.5 ml tube and 100 µl of 20% sarkosyl were added. After 2 min centrifugation, 600 µl were transferred to a QIAshredder spin column (Qiagen) placed in a 2 ml collection tube. The subsequent steps of RNA extraction were according to the manufacturer's protocol. The extracted RNA was then retro-transcribed into cDNA using the iScript cDNA Synthesis kit (Biorad, USA). For each sample, 2 µl of cDNA were amplified in a total volume of 20 µl containing 1X HotMaster Buffer, 0.5 µg/µl BSA and 1 U of HotMaster Taq DNA Polymerase (Eppendorf, Germany). Primers and RT-PCR parameters were chosen following the protocols reported in Table 1. 18S

Table 1 - List of references for primers and RT-PCR conditions

Target	RT-PCR assay
<i>Comoviridae</i>	
ArMV	Faggioli <i>et al.</i> , 2005
BBWV-1	Ferrer <i>et al.</i> , 2008
BBWV-2	Ferrer <i>et al.</i> , 2008
<i>Betaflexiviridae</i>	
AICV	Spence <i>et al.</i> , 2000
LSV	Lim <i>et al.</i> , 2009
<i>Bromoviridae</i>	
CMV	Faggioli <i>et al.</i> , 2005
<i>Bunyaviridae</i>	
IYSV	Kritzman <i>et al.</i> , 2000
INSV	Liu <i>et al.</i> , 2009
TSWV	Mumford <i>et al.</i> , 1994
<i>Potyviridae</i>	
AIMV	Spence <i>et al.</i> , 2000
BYMV	Ganesh Selvaraj <i>et al.</i> , 2009
DsMV	Wen-Chi <i>et al.</i> , 2010
KoMV	Wen-Chi <i>et al.</i> , 2010
LMoV	Lim <i>et al.</i> , 2009
TuMV	Wen-Chi <i>et al.</i> , 2010
ZaMV	Kwon <i>et al.</i> , 2003
ZaMMV	Wen-Chi <i>et al.</i> , 2010
<i>Tombusviridae</i>	
CarnMV	Cevik <i>et al.</i> , 2010
<i>Unassigned</i>	
TRV	Wei <i>et al.</i> , 2009

rRNA was used as internal control (Osman and Rowhani, 2006). Finally, 10 µl of the amplification mix was electrophoresed in a 1.5% agarose gel in TAE buffer [40 mM Tris base, 20 mM sodium acetate, 1 mM EDTA pH (8.0)]. The amplified cDNA fragments were visualized on a UV transilluminator.

Data were analyzed using Sigma-Plot software (version 11; Systat Software, San Jose, CA). The software was used to perform analysis of variance (ANOVA). Data expressed in percent were converted to arcsin values.  $P < 0.05$  was considered to be significant.

### 3. Results

The health status of Calla and Peruvian lily as determined by the present survey is reported in Table 2. *Z. aethiopica* was mainly affected by viruses belonging to the *Potyviridae* family. More than two plants out of three were infected by DsMV and 50%

Table 2 - Health status of Calla lily (*Zantedeschia aethiopica*) and Peruvian lily (*Alstroemeria* spp.) expressed as percentage of infected plants

Target	<i>Zantedeschia aethiopica</i>	<i>Alstroemeria</i> spp.
<i>Comoviridae</i>		
ArMV	-	-
BBWV-1	-	-
BBWV-2	-	-
<i>Betaflexiviridae</i>		
AICV	-	-
LSV	-	56.3 a
<i>Bromoviridae</i>		
CMV	-	12.5 b
<i>Bunyaviridae</i>		
IYSV	-	-
INSV	3.3 c *	-
TSWV	3.3 c	-
<i>Potyviridae</i>		
AIMV	-	-
DsMV	66.7 a	13.5 b
KoMV	-	-
LMoV	16.7 b	12.5 b
TuMV	-	-
ZaMMV	50.0 a	6.3 c
<i>Tombusviridae</i>		
CarnMV	-	-
<i>Unassigned</i>		
TRV	-	-

\* Values in the same column followed by the same letter do not differ significantly according to Duncan's multiple range test ( $P=0.05$ ).

Table 3 - Sequence of isolates of Zantedeschia mild mosaic virus and Lily mottle virus isolate detected in Italy

Zantedeschia mild mosaic virus isolate 2079 polyprotein gene, partial cds (GenBank: KF156666)

TCATTGAGTACCAACCCCAACAGTCCGATCTGTTAATACTCGCGCTCACAAACCAATTCAATAATTGGTATGATGCGATCAAAAATGAG-TATGGGGTTGATGATAGTCAGATGCAGAGAATCATGAATGGCTTCATGGTGTGGTGTCTCGAGAATGGGACATCACCAACATAAATGGCGTGTGGGT-TATGATGGATGGGGATGAACAAGTAGAATTTCCACTAAACCAATGGTGAGAAATGCCAAGCCTACGCTGCGTCAAATAATGCACCACTTTTCAGACG-CAGCCGAGGCTTACATTGAACCTTAGGAATGCCGCTGCCCATATATGCCTAGATATGGGTTGCTGCGGAACCTAAGAGACAGAGGTCTAGCACGCTTCG-CATTGACTTCTATGAAGTCACTTCAAAGACACCAGATCGTGCTAGAGAAGCTGTAGCGCAGATGAAGGCAGCAGCGCTAAACAATGTTTCCACAAG-GATGTTTGGATTGGATGGAAATATTGCAACTGCCACGGAGAACACTGAAAGGCACACTGCTAAGGATGTAAGTCCGAGCATGCACTCGCTACTCGG-GATCTCAGCCTTGCAAGTAAGGAGCTGGAAACAGCCACAGTTATTGTCTTGGATAGGGTTTAAATAGCCGTACTATTGTGCTTGTAGATGTTG-CAGTGTGGGCTCCACCTAAGGTTTATCAGTGTGGCTTTCCACCTAGTTCCTTACATTGCGCATAGTATGTG

Lily mottle virus isolate 2409 coat protein gene, partial cds (GenBank: KF156662.1)

TGCTGGGGCCTCTAGCTCCACACAAACGAGTCGCCAACACGTCCAGAGATTGCCGCGGTGATGTAGCACCACAACAGAGCTCTGAGGCTA-GAGTGGGTGATCGTGATGTTGATGCTGGCACCGTGGGAACATACCAAATCCCTCGACTGAAAGCACTGGCAACAAAGATTAACGTACCCAAGGT-CAAGGGGCGAACAATAGTGAACACTGGGCACCTTGTGAACATAACCCAGACCAACAGATATTTCAAATACAAGGTCAACCCAGAAGCAATTTGA-GACCTGGCATAACGCTGTGAAAGATGAGTATGGTCTCAACGACGAGAGTATGGCTCTCGCAATGAATGGTCTGATGGTTTGGTGCATAGAGAATGG-CACCTCACCAAAACATAAATGGCGTGTGGCTCATGATGACGAGATCAGCAAGTTGAATTTCTTTACGTCCTATCTTGAACACGCAAAACC-GACGCTGCGCAAAATTATGGCGCATTTCTCAAACCTCGCTGAAGCTTATATTGAGAAGCAAAATTTGGAGAAACCGTACATGCCTAGGTACGGCCTT-CAGCGAAATCTACCGATTTCAATCTAGCACGATTTGCTTTGATTCTATGA

of tested plants were infected by ZaMMV. Lower infection rates were reported for viruses belonging to *Bunyaviridae*. Even if *Alstroemeria* spp. plants were affected by *Potyviridae* viruses too, higher infection rates were recorded for *Betaflexiviridae*, where LSV infected more than half of plants. Further infections were caused by *Bromoviridae* viruses. *Comoviridae*, *Tombusviridae* or TRV infections were not found in both plants.

In *Z. aethiopica*, mixed infections were set at 40% for DsMV/ZaMMV, 6.7% for DsMV/LMoV/ZaMMV and 3.3% for DsMV/LMoV (data not shown). In *Alstroemeria* spp., mixed infections were set at 6.3% for LSV/ZaMMV, LSV/LMoV, DsMV/ZaMMV, CMV/LSV or CMV/DsMV/LSV.

With regard to *Alstroemeria* spp., the sequence obtained from a ZaMMV amplicon (GenBank accession no. KF156666 (Table 3) had 99% nucleotide identity with the corresponding fragment of a reference ZaMMV isolate (GenBank accession no. AY626825). Further isolates of LMoV were detected (GenBank accessions no. KF156667, KF156668, KF156669). Each further isolate had 99% nucleotide identity with the corresponding fragment of a reference ZaMMV isolate. All four isolates are different but had 99% nucleotide identity.

The sequence obtained from LMoV amplicon (GenBank accession no. KF156662) (Table 3) had 94% nucleotide identity with the corresponding fragment of a reference LMoV isolate (GenBank accession no. JN703466). The same LMoV amplicon was obtained from a *Z. aethiopica* sample as well. Further isolates of LMoV were detected in both species with 100% nucleotide identity with KF156662 (GenBank accessions no. KF156663, KF156664, KF156665).

#### 4. Conclusions

Various viral infections, mainly due to viruses belonging to two families, *Betaflexiviridae* and *Potyviridae*, seem to affect Calla and Peruvian lily in Tuscan nurseries. Most detected viruses are frequently reported for both plants and mixed infection of virus belonging to *Potyviridae* are quite common in *Z. aethiopica* (Huang et al., 2007). However, some evidence is rarer.

To our knowledge this is the first report of LMoV in *Alstroemeria* spp. and *Z. aethiopica* or ZaMMV in *Alstroemeria* spp. in Italy, while ZaMMV was recently identified in Taiwan (Huang and Chang, 2005) and in Italy (Rizzo et al., 2015 b).

This report puts in evidence how widespread the virus is within these common plants and the need for constant monitoring of the health status for flower production. Even if some of viruses that affect Calla and Peruvian lily may be eradicated by thermotherapy (Panattoni et al., 2013) and heat treatment may help in *Bromoviridae* control (Luvisi et al., 2015), prevention represents the preferred method of virus control.

#### References

- CEVIK B., BAKIR T., KOCA G., 2010 - First report of Carnation mottle virus in Turkey. - Plant Pathol., 59: 394.
- FAGGIOLI F., FERRETTI L., ALBANESE G., SCIARRONI R., PASQUINI G., LUMIA V., BARBA M., 2005 - Distribution of olive tree viruses in Italy as revealed by one-step RT-PCR. - J. Plant Pathol., 87: 49-55.

- FERRER R., ESCRIBU F., LUIS-ARTEAGA M., GUERRI J., MORENO P., RUBIO L., 2008 - *New molecular methods for identification of Broad bean wilt virus 1*. - Mol. Cell. Probes, 22: 223-227.
- GANESH SELVARAJ D., POKORNY R., HOLKOVA L., 2009 - *Comparative analysis of ELISA, one step RT-PCR and IC-RT-PCR for the detection of bean yellow Mosaic virus in gladiolus*. - Commun. Agric. Appl. Biol. Sci., 74: 853-859.
- HUANG C.H., CHANG Y.C., 2005 - *Identification and molecular characterization of Zantedeschia mild mosaic virus, a new calla-lily-infecting potyvirus*. - Arch. Virol., 150: 1221-1230.
- HUANG C.H., HU W.C., YANG T.C., CHANG Y.C., 2007 - *Zantedeschia mild mosaic virus, a new widespread virus in calla lily, detected by ELISA, dot-blot hybridization and IC-RT-PCR*. - Plant Pathol., 56: 183-189.
- KRITZMAN A., BECKELMAN H., ALEXANDROV S., COHEN J., LAMPEL M., ZEIDAN M., RACCAH B., GERA A., 2000 - *Lisianthus leaf necrosis: A new disease of Lisianthus caused by Iris yellow spot virus*. - Plan Dis., 84: 1185-1189.
- KWON S.B., HA J.H., YOON J.Y., RYU K.H., 2003 - *Zantedeschia mosaic virus causing leaf mosaic symptom in calla lily is a new potyvirus*. - Arch. Virol., 147: 2281-2289.
- LIM H.J., BAE E.H., LEE Y.J., PARK S.H., LEE K.J., KIM S.R., 2009 - *Detection of Lily symptomless virus, Lily mottle virus, and Cucumber mosaic virus from Lilium grown in Korea by RT-PCR*. - Korean Journal of Microbiology, 45: 251-256.
- LIU H.Y., SEARS J.L., MOU B., 2009 - *Spinach (Spinacia oleracea) is a new natural host of impatiens necrotic spot virus in California*. - Plant Dis., 93: 673.
- LUVISI A., PANATTONI A., MATERAZZI A., 2015 - *Heat treatments for sustainable control of soil viruses*. - Agron. Sustain. Dev., 35: 657-666.
- MACKENZIE D.J., MCLEAN M.A., MUKERJI S., GREEN M., 1997 - *Improved RNA extraction from woody plants for the detection of viral pathogens by reverse transcription-polymerase chain reaction*. - Plant Dis., 81: 222-226.
- MUMFORD R.A., BARKER I., WOOD K.R., 1994 - *The detection of tomato spotted wilt virus using the polymerase chain reaction*. - J. Virol. Methods, 46: 303-311.
- OSMAN F., ROWHANI A., 2006 - *Application of a spotting sample preparation technique for the detection of pathogens in woody plants by RT-PCR and real-time PCR (TaqMan)*. - J. Virol. Methods, 133: 130-136.
- PANATTONI A., LUVISI A., TRIOLO E., 2013 - *Elimination of viruses in plants: twenty years of progress*. - Span. J. Agric. Res., 11: 173-188.
- PARK T.H., HAN I.S., KIM J.B., 2010 - *Review on the development of virus resistant plants in Alstroemeria*. - J. Plant Biotechnol., 37: 370-378.
- RIZZO D., MATERAZZI A., STEFANI L., FARINA P., VANARELLI S., PANATTONI A., LUVISI A., 2015 a - *Distribution of regulated viruses in cv. Sangiovese vineyards in Tuscany*. - J. Plant Pathol., 97: 131-135.
- RIZZO D., PANATTONI A., STEFANI L., PAOLI M., NESI B., LAZZERESCHI S., VANARELLI S., FARINA P., DELLA BARTOLA M., MATERAZZI A., LUVISI A., 2015 b - *First report of Zantedeschia mild mosaic virus on Zantedeschia Aethiopica (L) Spreng in Italy*. - J. Plant Pathol., 92: 399.
- RIZZO D., STEFANI L., PAOLI M., TRIOLO E., PANATTONI A., LUVISI A., 2012 - *The sustainability of old grapevine mother plants in relation to new mandatory diagnostic tests for virus control*. - Adv. Hort. Sci., 26(3-4): 148-150.
- SPENCE N.J., MILLS P.R., BARBARA D.J., 2000 - *A survey of viruses of Alstroemeria in the UK and the characterization of carlaviruses infecting Alstroemeria*. - Eur. J. Plant Pathol., 106: 843-847.
- WEI T., LU G., CLOVER G.R.G., 2009 - *A multiplex RT-PCR for the detection of Potato yellow vein virus, Tobacco rattle virus and Tomato infectious chlorosis virus in potato with a plant internal amplification control*. - Plant Pathol., 58: 203-209.
- WEN-CHI H., CHIN-HSING H., SHU-CHUAN L., CHUN-I W., YA-CHUN C., 2010 - *Detection of four calla potyviruses by multiplex RT-PCR using nad5 mRNA as an internal control*. - Eur. J. Plant Pathol., 126: 43-52.